

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number
WO 01/49664 A1

- (51) International Patent Classification⁷: **C07D 207/34**, 401/12, 409/14, 407/12, 413/12, 409/12, A01N 43/36
- (74) Agent: **BASTIAN, Werner**; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).
- (21) International Application Number: PCT/EP00/11196
- (22) International Filing Date:
11 November 2000 (11.11.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
9930750.6 29 December 1999 (29.12.1999) GB
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **SYNGENTA PARTICIPATIONS AG** [CH/CH]; Schwabzaldallee 215, CH-4058 Basel (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **WALTER, Harald** [DE/CH]; Chilchmattstrasse 12b, CH-4118 Rodersdorf (CH). **TRAH, Stephan** [DE/DE]; Andreas Hofer-Strasse 150a, 79111 Freiburg im Breisgau (DE). **SCHNEIDER, Hermann** [DE/DE]; Alemannenstrasse 17, 79423 Heitersheim (DE).
- Published:**
— With international search report.
— With amended claims.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TRIFLUOROMETHYLPYRROLE CARBOXAMIDES AND TRIFLUOROMETHYLPYRROLETHIOAMIDES AS FUNGICIDES




(57) Abstract: Novel pyrrole derivatives of formula (I) wherein X is oxygen or sulfur; R₁ is hydrogen, C₁-C₄alkyl unsubstituted or substituted, or halogen; R₂ is C₁-C₄alkyl unsubstituted or substituted; and A is orthosubstituted aryl; orthosubstituted heteroaryl; bicycloaryl unsubstituted or substituted; or bicycloheteroaryl unsubstituted or substituted. The novel compounds have plant-protective properties and are suitable for protecting plants against infestations by phytopathogenic microorganisms.

Trifluoromethylpyrrolecarboxamides and trifluoromethylpyrrolethioamides as fungicides

The present invention relates to novel trifluoromethylpyrrolecarboxamides or trifluoromethylpyrrolethioamides which have microbicidal activity, in particular fungicidal activity. The invention also relates to the preparation of these substances, to agrochemical compositions which comprise at least one of the novel compounds as active ingredient, to the preparation of the compositions mentioned and to the use of the active ingredients or compositions in agriculture and horticulture for controlling or preventing infestation of plants by phytopathogenic microorganisms, preferably fungi.

The trifluoromethylpyrrolecarboxamides of the present invention have the general formula I



wherein

X is oxygen or sulfur;

R₁ is hydrogen, C₁-C₄alkyl unsubstituted or substituted, C₁-C₄alkoxy unsubstituted or substituted, or halogen;

R₂ is C₁-C₄alkyl unsubstituted or substituted; and

A is orthosubstituted aryl; orthosubstituted heteroaryl; bicycloaryl unsubstituted or substituted; or bicycloheteroaryl unsubstituted or substituted.

Surprisingly, it has now been found that the compounds of formula I exhibit improved biological properties which render them more suitable for the practical use in agriculture and horticulture.

Where asymmetrical carbon atoms are present in the compounds of formula I, these compounds are in optically active form. The invention relates to the pure isomers, such as enantiomers and diastereomers, as well as to all possible mixtures of isomers, e.g. mixtures of diastereomers, racemates or mixture of racemates.

Within the present specification alkyl denotes methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl and tert-butyl. Non-branched alkyl is preferred. Alkyl as part of other radicals such as alkoxy, haloalkyl, alkylcycloalkyl, alkylcycloalkoxy, etc. is understood in an analogous way. Halogen will be understood generally as meaning fluoro, chloro, bromo or iodo. Fluoro, chloro or bromo are preferred meanings. Halogen as part of other radicals such as haloalkyl, haloalkoxy, haloalkenyl, haloalkenyloxy, haloaryl or haloheteroaryl, etc. is understood in an analogous way. Haloaryl or haloheteroaryl designates mono- to five times halo-substituted aryl, whereby the halogens are independently chosen. Where more than two halogens are present, the halogens are preferably the same, e.g. trifluorophenyl, trichlorophenyl, tetrachlorophenyl or perchlorophenyl.

Aryl is phenyl or naphthyl.

Heteroaryl will be understood as a 5- to 10-membered ring that may contain up to 3 heteroatoms, such as nitrogen, oxygen or sulfur. The following list of examples is not exhaustive : furyl, thienyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, pyrrolyl, dithiolyl, oxathiolyl, dioxazolyl, oxathiazolyl, oxathiolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl, benzofuranyl, benzimidazolyl, indazolyl, benzotriazolyl, benzothiazolyl, benzoxazolyl, quinolinyl, isoquinolinyl, phthalazinyl, quinoxalinyl, quinazolinyl, cinnolinyl, naphthyridinyl, isobenzofuranyl, isoindolyl, benzothiadiazolyl, benzothienyl, benzisoxazolyl, purinyl, 5,6-dihydro-1,4,2-dioxazinyl, and the like.

Bicycloaryl or bicycloheteroaryl will be understood as a 6-membered aryl or 6-membered heteroaryl, wherein it may contain up to 3 heteroatoms such as nitrogen, oxygen or sulfur, and which is fused to an additional ring. The fused ring may be aromatic, partially hydrogenated or completely saturated, may be a ring from 5 to 7 ring members, of which up to 3 members may be heteroatoms selected from the group nitrogen, oxygen and sulfur. The following list of examples is not exhaustive: dihydroisobenzofuranyl, dihydroisoindolyl, and the like.

Preferred embodiments of compounds of the formula I are those wherein X is oxygen or sulfur; or

X is oxygen; or

X is sulfur; or

R₁ is hydrogen, C₁-C₄alkyl unsubstituted or substituted, C₁-C₄alkoxy unsubstituted or substituted, or halogen; or

R₁ is hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄haloalkoxy-C₁-C₄alkyl or halogen; or

R₁ is hydrogen, C₁-C₄alkyl or C₁-C₄haloalkyl; or

R₁ is hydrogen or C₁-C₃alkyl; or

R₂ is C₁-C₄alkyl unsubstituted or substituted; or


R₂ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy-C₁-C₄alkyl or C₁-C₄haloalkoxy-C₁-C₄alkyl; or


R₂ is C₁-C₄alkyl, C₁-C₃haloalkyl or C₁-C₃alkoxy-C₁-C₃alkyl; or

R₂ is C₁-C₃alkyl or C₁-C₃alkoxy-C₁-C₃alkyl; or

A is orthosubstituted aryl; orthosubstituted heteroaryl; bicycloaryl unsubstituted or substituted; or bicycloheteroaryl unsubstituted or substituted; or

A is a group





A is A1, A2, A3, A5, A8, A10, A13, A14, A17, A18, A20, A21, A22, A24, A25, A26, A27, A29, A31 or A32; or

A is A1, A2, A3, A17, A20, A21, A24, A25, A26, A27 or A31; or

R₃ is C₃-C₇cycloalkyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₁-C₄alkoxy, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkoxy or C₁-C₄alkyl; C₄-C₇cycloalkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₁-C₄alkoxy, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkoxy or C₁-C₄alkyl; C₆-C₇cyclodialkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₁-C₄alkoxy, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkoxy or C₁-C₄alkyl; thienyl, furyl, pyrrolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, thiadiazolyl, imidazolyl, tetrazolyl, triazinyl, benzothienyl, 5,6-dihydro-1,4,2-dioxazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl, which are unsubstituted or substituted by halogen, C₁-C₆haloalkyl, C₁-C₆alkyl, hydroxy, cyano, nitro, CHO, C₁-C₄alkoxy-C₁-C₄alkyl, COOC₁-C₄alkyl, C₁-C₆haloalkoxy-C₁-C₄alkyl, C₁-C₆alkoxy or C₁-C₆haloalkoxy; or

R₃ is C₅-C₇cycloalkyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkoxy or C₁-C₄alkyl; C₅-C₇cycloalkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₁-C₃alkoxy, C₂-C₄alkenyl, C₂-C₄alkynyl or

C₁-C₄alkyl; C₆-C₇cyclodialkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₁-C₃alkoxy, C₂-C₄alkenyl, C₂-C₄alkynyl or C₁-C₃alkyl; thienyl, furyl, oxazolyl, isoxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl, which are unsubstituted or substituted by halogen, C₁-C₄haloalkyl, C₁-C₄alkyl, hydroxy, C₁-C₄alkoxy or C₁-C₄haloalkoxy; or

R₃ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or mono- to disubstituted by chloro, bromo, C₁-C₂alkyl, C₁-C₂haloalkyl or C₁-C₂haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; or

R₃₁ is C₃-C₇cycloalkyl, C₃-C₇cycloalkenyl or C₅-C₇cycloalkadienyl which are unsubstituted or substituted by halogen, C₁-C₆alkoxy, C₁-C₆alkoxy-C₁-C₆alkyl, C₁-C₆haloalkoxy-C₁-C₄alkyl, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄haloalkoxy, C₂-C₄alkenyl or C₂-C₅alkynyl; phenyl unsubstituted or substituted by halogen, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₅alkynyl, CHO, COOC₁-C₄alkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₄alkyl-C₁-C₄alkoxy, C₁-C₄haloalkoxy-C₁-C₄alkyl, C₁-C₄haloalkyl-C₁-C₄alkoxy, cyano or nitro; thienyl, furyl, pyrrolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, thiadiazolyl, imidazolyl, triazinyl, pyridyl, pyridazinyl, pyrazinyl or pyrimidinyl, which are unsubstituted or substituted by halogen, C₁-C₆haloalkyl, C₁-C₆alkyl, hydroxy, cyano, nitro, CHO, COOC₁-C₄alkyl, C₁-C₆haloalkoxy-C₁-C₄alkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₆alkoxy or C₁-C₆haloalkoxy; or

R₃₁ is C₅-C₇cycloalkyl, C₅-C₇cycloalkenyl or C₅-C₇cycloalkadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; phenyl which is unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; or

R₃₁ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; or

R₄ is hydrogen; cyano; nitro; halogen; C₁-C₄alkoxy; C₁-C₄haloalkyl; C₁-C₄alkyl; C₁-C₄alkoxy-C₁-C₄alkyl; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₄haloalkoxy-C₁-C₄alkyl; or C₁-C₄haloalkoxy; or

R_4 is hydrogen; halogen; C_1 - C_4 alkoxy; C_1 - C_4 alkyl; C_5 - C_7 cycloalkyl unsubstituted or substituted by C_1 - C_3 alkyl or C_1 - C_3 haloalkyl; C_1 - C_4 haloalkyl; or C_1 - C_4 haloalkoxy; or

R_4 is hydrogen; halogen; C_1 - C_3 alkoxy; C_1 - C_3 haloalkyl; C_1 - C_3 alkyl; C_5 - C_7 cycloalkyl unsubstituted or substituted by C_1 - C_3 alkyl or C_1 - C_3 haloalkyl; or C_1 - C_3 haloalkoxy; or

R_5 , R_6 , R_7 , R_8 and R_9 are identical or different and are each independently of the others hydrogen, halogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkyl, C_1 - C_6 haloalkoxy- C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 alkoxy- C_1 - C_4 alkyl or C_3 - C_7 cycloalkyl; or

R_5 , R_6 , R_7 and R_8 are identical or different and are each independently of the others hydrogen, C_1 - C_4 haloalkyl, C_1 - C_4 alkyl, C_1 - C_4 alkoxy or C_1 - C_4 haloalkoxy; or


R_5 , R_6 , R_7 and R_8 are identical or different and are each independently of the others hydrogen or C_1 - C_3 alkyl.


Within the group of compounds of formula I those compounds are preferred wherein :

R_1 is hydrogen, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy- C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkoxy, C_1 - C_4 haloalkoxy- C_1 - C_4 alkyl or halogen ;

R_2 is C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 alkoxy- C_1 - C_4 alkyl or C_1 - C_4 haloalkoxy- C_1 - C_4 alkyl ;

A is a group





and

R_3 is C_3 - C_7 cycloalkyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkoxy or C_1 - C_4 alkyl; C_4 - C_7 cycloalkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkoxy or C_1 - C_4 alkyl; C_6 - C_7 cyclodialkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkoxy or C_1 - C_4 alkyl; thienyl, furyl, pyrrolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, thiadiazolyl, imidazolyl, tetrazolyl, triazinyl, benzothienyl, 5,6-dihydro-1,4,2-dioxazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl, which are unsubstituted or substituted by halogen, C_1 - C_6 haloalkyl, C_1 - C_6 alkyl, hydroxy, cyano, nitro, CHO, C_1 - C_4 alkoxy- C_1 - C_4 alkyl, COOC- C_4 alkyl, C_1 - C_6 haloalkoxy- C_1 - C_4 alkyl, C_1 - C_6 alkoxy or C_1 - C_6 haloalkoxy;

R_{31} is C_3 - C_7 cycloalkyl, C_3 - C_7 cycloalkenyl or C_5 - C_7 cycloalkadienyl which are unsubstituted or substituted by halogen, C_1 - C_6 alkoxy, C_1 - C_6 alkoxy- C_1 - C_6 alkyl, C_1 - C_6 haloalkoxy- C_1 - C_4 alkyl, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, C_2 - C_4 alkenyl or C_2 - C_5 alkynyl; phenyl unsubstituted or substituted by halogen, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, C_1 - C_4 alkyl, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_5 alkynyl, CHO, COOC- C_4 alkyl, C_1 - C_4 alkoxy- C_1 - C_4 alkyl,

C₁-C₄alkyl-C₁-C₄alkoxy, C₁-C₄haloalkoxy-C₁-C₄alkyl, C₁-C₄haloalkyl-C₁-C₄alkoxy, cyano or nitro; thienyl, furyl, pyrrolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, thiadiazolyl, imidazolyl, triazinyl, pyridyl, pyridazinyl, pyrazinyl or pyrimidinyl, which are unsubstituted or substituted by halogen, C₁-C₆haloalkyl, C₁-C₆alkyl, hydroxy, cyano, nitro, CHO, COOC₁-C₄alkyl, C₁-C₆haloalkoxy-C₁-C₄alkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₆alkoxy or C₁-C₆haloalkoxy;

R₄ is hydrogen; cyano; nitro; halogen; C₁-C₄alkoxy; C₁-C₄haloalkyl; C₁-C₄alkyl; C₁-C₄alkoxy-C₁-C₄alkyl; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₄haloalkoxy-C₁-C₄alkyl; or C₁-C₄haloalkoxy; and

R₅, R₆, R₇, R₈ and R₉ are identical or different and are each independently of the others hydrogen, halogen, C₁-C₄haloalkyl, C₁-C₄alkyl, C₁-C₆haloalkoxy-C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkoxy-C₁-C₄alkyl or C₃-C₇cycloalkyl (subgroup AA).

Cycloalkyl is, depending on the ring size, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

Cycloalkenyl is, depending on the ring size, cyclopentenyl, cyclohexenyl or cycloheptenyl.

Cycloalkadienyl is, depending on the ring size, cyclopentadienyl, cyclohexadienyl or cycloheptadienyl.

Alkenyl will be understood as meaning straight-chain or branched alkenyl such as allyl, methallyl, 1-methylvinyl or but-2-en-1-yl. Preferred alkenyl radicals contain 3 to 4 carbon atoms in the chain.

Alkynyl can likewise, in accordance with the number of carbon atoms, be straight-chain or branched and is typically propargyl, but-1-yn-1-yl or but-1-yn-3yl.

Within the group AA of compounds of formula I those compounds are preferred wherein X is oxygen (subgroup AB).

Another group of compounds of formula I within the group AA are those wherein X is sulfur (subgroup AC).

Within the subgroup AB are those compounds preferred wherein

R₁ is hydrogen, C₁-C₄alkyl or C₁-C₄haloalkyl;

R₂ is C₁-C₄alkyl, C₁-C₃haloalkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

A is A1, A2, A3, A5, A8, A10, A13, A14, A17, A18, A20, A21, A22, A24, A25, A26, A27, A29, A31 or A32;

R₃ is C₅-C₇cycloalkyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₅-C₇cycloalkenyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₆-C₇cyclodialkenyl, unsubstituted or mono- to disubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl, which are unsubstituted or substituted by halogen, hydroxy, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy or C₁-C₄haloalkoxy;

R₃₁ is C₅-C₇cycloalkyl, C₅-C₇cycloalkenyl or C₅-C₇cycloakadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; phenyl which is unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₄ is hydrogen; halogen; C₁-C₄alkyl; C₁-C₄alkoxy; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₄haloalkyl; or C₁-C₄haloalkoxy; and

R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy (subgroup AB1).

Within the subgroup AB1 are those compounds more preferred wherein

A is A1, A2, A3, A17, A20, A21, A24, A25, A26, A27 or A31;

R₁ is hydrogen or C₁-C₃alkyl;

R₂ is C₁-C₃alkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

R₃ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or mono- to disubstituted by chloro, bromo, C₁-C₂alkyl, C₁-C₂haloalkyl or C₁-C₂haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₃₁ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, triazinyl,

pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₄ is hydrogen; halogen; C₁-C₃alkyl; C₁-C₃haloalkyl; C₁-C₃alkoxy; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; or C₁-C₃haloalkoxy; and R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen or C₁-C₃alkyl (subgroup AB2).

Within the subgroup AC are those compounds preferred wherein

R₁ is hydrogen, C₁-C₄alkyl or C₁-C₄haloalkyl;

R₂ is C₁-C₄alkyl, C₁-C₃haloalkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

A is A1, A2, A3, A5, A8, A10, A13, A14, A17, A18, A20, A21, A22, A24, A25, A26, A27, A29, A31 or A32;

R₃ is C₅-C₇cycloalkyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₅-C₇cycloalkenyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₆-C₇cyclodialkenyl, unsubstituted or mono- to disubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl, which are unsubstituted or substituted by halogen, hydroxy, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy or C₁-C₄haloalkoxy;

R₃₁ is C₅-C₇cycloalkyl, C₅-C₇cycloalkenyl or C₅-C₇cycloakadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; phenyl which is unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; R₄ is hydrogen; halogen; C₁-C₄alkyl; C₁-C₄alkoxy; C₁-C₄haloalkyl; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; or C₁-C₄haloalkoxy; and R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy (subgroup AC1).

Within the subgroup AC1 are those compounds more preferred wherein

A is A1, A2, A3, A17, A20, A21, A24, A25, A26, A27 or A31;

R₁ is hydrogen or C₁-C₃alkyl;

R₂ is C₁-C₃alkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

R₃ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or mono- to disubstituted by chloro, bromo, C₁-C₂alkyl, C₁-C₂haloalkyl or C₁-C₂haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₃₁ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₄ is hydrogen; halogen; C₁-C₃alkyl; C₁-C₃haloalkyl; C₁-C₃alkoxy; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; or C₁-C₃haloalkoxy; and

R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen or C₁-C₃alkyl (subgroup AC2).

Within the subgroup AB are those compounds preferred wherein

A is A1 and

R₃ is C₅-C₇cycloalkyl unsubstituted or monosubstituted by C₁-C₄alkyl or C₁-C₄haloalkyl (subgroup AB3).

Preferred individual compounds are :

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-methylcyclohexyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-ethylcyclohexyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-trifluoromethylcyclohexyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-methylcyclopentyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-ethylcyclopentyl)-2-benzamide];


1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-trifluoromethylcyclopentyl)-2-benzamide].

The compounds according to formula I may be prepared according to the following reaction schemes.

A) Synthesis of the pyrrole carboxylic acids :


Route 1 (Tosmic-route)


Scheme 1



Route 2 (Trifluoroacetoacetic acid-route, analogous to JP-07157466)

Scheme 2






Base 1 = $NaHCO_3$, Na_2CO_3 , $KHCO_3$, K_2CO_3 , $CaCO_3$ and other bases

Base 2 = $NaOH$, KOH , NaH , KH , $n-BuLi$ and others

B) Synthesis of the amides/thioamides


Scheme 3



Base = NEt_3 , Hünig-base, Na_2CO_3 , K_2CO_3 and others


B1) Synthesis of the amides wherein A is phenyl, pyridine or pyrimidine

Scheme 4



"Pd-catalyst"-system
solvent, base

$R_3-B(OH)_2$
(Suzuki-coupling)



G = phenyl, pyridine, pyrimidine

The "in situ" prepared pyrazole carboxylic acid chloride reacts with an ortho-halosubstituted phenylamine in the presence of a solvent like toluene, benzene, xylene, hexane, cyclohexane, THF, chloroform or methylenechloride and in the presence of a base like sodium carbonate, sodium hydrogencarbonate, potassium carbonate, Hünig base, triethylamine or pyridine at a temperature between 0°C and reflux temperature. The obtained pyrazolecarboxamide reacts with a boronic acid of the formula $R_3-B(OH)_2$ in the presence of a Pd-catalyst like $Pd(P(phenyl)_3)_4$, $Pd(P(phenyl)_3)_2Cl_2$, $PdCl_2dppb$, $Pd_2(dba)_3$, $Pd(OAc)_2$, $PdOAc_2/(o-tolyl)_3P$, $Pd(OAc)_2/dppf$, $Pd(PhCN)_2Cl_2/Ph_3As$, $Pd(CH_3CN)_2Cl_2$, $Pd_2(dba)_3/P(tert.butyl)_3$, $Pd(OAc)_2/P(tert.butyl)_2biphenyl$, $Pd(OAc)_2/TPPTS$, $Pd(OAc)_2/PCy_3$, $Pd(OAc)_2/P(O-i-Pr)_3$, $Pd(OAc)_2/2$ -dimethylamino-2'-dicyclohexylphosphinobiphenyl, $Pd(OAc)_2/2$ -dimethylamino-2'-ditert.butylphosphinobiphenyl, $Pd(OAc)_2/(o-biphenyl)P(cyclohexyl)_2$ in a solvent like 1,2-dimethoxyethane/water, DMF, DMA, THF/water, dioxane/water, benzene, toluene, xylene and others and a base like sodium carbonate, sodium hydrogencarbonate, potassium carbonate, cesium carbonate, potassium phosphate, triethylamine, sodium hydroxide, sodium ethylate, sodium tert.butylate, silver oxide, barium carbonate, potassium fluoride or cesium fluoride at a temperature between 0°C and reflux temperature.

C) Synthesis of the ortho-substituted amines A-NH₂ :

The compounds are either known from the literature or can be prepared by known methods. For example the following amines or important intermediates for the synthesis of the amines can be prepared according to the following literature:

A₁-NH₂ : Tetrahedron 1993,49,49-64 or EP-83975 or J.Org.Chem. 1995,60,292;

A₂-NH₂ : EP-737682;

A₃-NH₂ : J.Chem.Res.(S) 1978,11,428 or Chem.Scr.1972,2,245;

A₁₇-NH₂ : Org.Prep.Procedures Int. 1989,21,141;
A₂₁-NH₂ : J.Chem.Soc.Perkin I 1981,5,1591;
A₂₅-NH₂ : Tetrahedron 1993,49,49-64 or Heterocycles 1999,51,721;
A₂₆-NH₂ : Synthesis 1996,10,1015 or Synthesis 1994,9,931;
A₂₇-NH₂ : Liebigs Ann.Chemie 1977,537-544 or J.Med.Chem. 1975,18,623;
A₃₁-NH₂ : EP-315502;
A₃₂-NH₂ : J.Pesticide Science 1993,18,245;
A₃₀-NH₂ : J.Heterocyclic Chem. 1982,19,1285.

Surprisingly, it has now been found that the novel compounds of formula I have, for practical purposes, a very advantageous spectrum of activities for protecting plants against diseases that are caused by fungi as well as by bacteria and viruses.

The compounds of formula I can be used in the agricultural sector and related fields of use as active ingredients for controlling plant pests. The novel compounds are distinguished by excellent activity at low rates of application, by being well tolerated by plants and by being environmentally safe. They have very useful curative, preventive and systemic properties and are used for protecting numerous cultivated plants. The compounds of formula I can be used to inhibit or destroy the pests that occur on plants or parts of plants (fruit, blossoms, leaves, stems, tubers, roots) of different crops of useful plants, while at the same time protecting also those parts of the plants that grow later e.g. from phytopathogenic microorganisms.

It is also possible to use compounds of formula I as dressing agents for the treatment of plant propagation material, in particular of seeds (fruit, tubers, grains) and plant cuttings (e.g. rice), for the protection against fungal infections as well as against phytopathogenic fungi occurring in the soil.

The compounds I are, for example, effective against the phytopathogenic fungi of the following classes: Fungi imperfecti (e.g. Botrytis, Pyricularia, Helminthosporium, Fusarium, Septoria, Cercospora and Alternaria) and Basidiomycetes (e.g. Rhizoctonia, Hemileia, Puccinia). Additionally, they are also effective against the Ascomycetes classes (e.g. Venturia and Erysiphe, Podosphaera, Monilinia, Uncinula) and of the Oomycetes classes (e.g. Phytophthora, Pythium, Plasmopara). Outstanding activity has been observed against

powdery mildew (*Erysiphe* spp.). Furthermore, the novel compounds of formula I are effective against phytopathogenic bacteria and viruses (e.g. against *Xanthomonas* spp, *Pseudomonas* spp, *Erwinia amylovora* as well as against the tobacco mosaic virus).

Within the scope of present invention, target crops to be protected typically comprise the following species of plants: cereal (wheat, barley, rye, oat, rice, maize, sorghum and related species); beet (sugar beet and fodder beet); pomes, drupes and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries); leguminous plants (beans, lentils, peas, soybeans); oil plants (rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans, groundnuts); cucumber plants (pumpkins, cucumbers, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, mandarins); vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, paprika); lauraceae (avocado, cinnamomum, camphor) or plants such as tobacco, nuts, coffee, eggplants, sugar cane, tea, pepper, vines, hops, bananas and natural rubber plants, as well as ornamentals.

The compounds of formula I are used in unmodified form or, preferably, together with the adjuvants conventionally employed in the art of formulation. To this end they are conveniently formulated in known manner to emulsifiable concentrates, coatable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granulates, and also encapsulations e.g. in polymeric substances. As with the type of the compositions, the methods of application, such as spraying, atomising, dusting, scattering, coating or pouring, are chosen in accordance with the intended objectives and the prevailing circumstances. The compositions may also contain further adjuvants such as stabilisers, antifoams, viscosity regulators, binders or tackifiers as well as fertilisers, micronutrient donors or other formulations for obtaining special effects.

Suitable carriers and adjuvants can be solid or liquid and are substances useful in formulation technology, e.g. natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilisers. Such carriers are for example described in WO 97/33890.

The compounds of formula I are normally used in the form of compositions and can be applied to the crop area or plant to be treated, simultaneously or in succession with further compounds. These further compounds can be e.g. fertilisers or micronutrient donors or other preparations which influence the growth of plants. They can also be selective herbicides as well as insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures of several of these preparations, if desired together with further carriers, surfactants or application promoting adjuvants customarily employed in the art of formulation.

The compounds of formula I can be mixed with other fungicides, resulting in some cases in unexpected synergistic activities.

Mixing components which are particularly preferred are azoles such as azaconazole, bitertanol, propiconazole, difenoconazole, diniconazole, cyproconazole, epoxiconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imazalil, imibenconazole, ipconazole, tebuconazole, tetraconazole, fenbuconazole, metconazole, myclobutanil, perfurazoate, penconazole, bromuconazole, pyrifenoxy, prochloraz, triadimefon, triadimenol, triflumizole or triticonazole; pyrimidinyl carbinols such as ancymidol, fenarimol or nuarimol; 2-amino-pyrimidine such as bupirimate, dimethirimol or ethirimol; morpholines such as dodemorph, fenpropidin, fenpropimorph, spiroxamin or tridemorph; anilinopyrimidines such as cyprodinil, pyrimethanil or mepanipyrim; pyrroles such as fenpiclonil or fludioxonil; phenylamides such as benalaxyl, furalaxyl, metalaxyl, R-metalaxyl, ofurace or oxadixyl; benzimidazoles such as benomyl, carbendazim, debacarb, fuberidazole or thiabendazole; dicarboximides such as chlozolinate, dichlozoline, iprodione, myclozoline, procymidone or vinclozolin; carboxamides such as carboxin, fenfuram, flutolanil, mepronil, oxycarboxin or thifluzamide; guanidines such as guazatine, dodine or iminoctadine; strobilurines such as azoxystrobin, kresoxim-methyl, metominostrobin, SSF-129, methyl 2-[(2-trifluoromethyl)-pyrid-6-yloxy-methyl]-3-methoxyacrylate or 2-[α [[α -methyl-3-trifluoromethyl-benzyl]imino]-oxy]-o-tolyl]-glyoxylic acid-methylester-O-methyloxime (trifloxystrobin); dithiocarbamates such as ferbam, mancozeb, maneb, metiram, propineb, thiram, zineb or ziram; N-halomethylthio-dicarboximides such as captan, captan, dichlofluanid, fluoromide, folpet or tolyfluanid; copper compounds such as Bordeaux mixture, copper hydroxide, copper oxychloride, copper sulfate, cuprous oxide, mancozeb or oxine-copper; nitrophenol derivatives such as dinocap or nitrothal-isopropyl; organo phosphorous derivatives such as edifenphos, iprobenphos, isoprothiolane, phosdiphen, pyrazophos or toclofos-methyl; and other compounds of diverse structures such

as acibenzolar-S-methyl, anilazine, blasticidin-S, chinomethionat, chloroneb, chlorothalonil, cymoxanil, dichlone, diclomezine, dicloran, diethofencarb, dimethomorph, dithianon, etridiazole, famoxadone, fenamidone, fentin, ferimzone, fluazinam, flusulfamide, fenhexamid, fosetyl-aluminium, hymexazol, kasugamycin, methasulfocarb, pencycuron, phthalide, polyoxins, probenazole, propamocarb, pyroquilon, quinoxifen, quintozone, sulfur, triazoxide, tricyclazole, triforine, validamycin, (S)-5-methyl-2-methylthio-5-phenyl-3-phenyl-amino-3,5-dihydroimidazol-4-one (RPA 407213), 3,5-dichloro-N-(3-chloro-1-ethyl-1-methyl-2-oxopropyl)-4-methylbenzamide (RH-7281), N-allyl-4,5-dimethyl-2-trimethylsilylthiophene-3-carboxamide (MON 65500), 4-chloro-4-cyano-N,N-dimethyl-5-p-tolylimidazole-1-sulfonamide (IKF-916), N-(1-cyano-1,2-dimethylpropyl)-2-(2,4-dichlorophenoxy)-propionamide (AC 382042), or iprovalicarb (SZX 722).

A preferred method of applying a compound of formula I, or an agrochemical composition which contains at least one of said compounds, is foliar application. The frequency of application and the rate of application will depend on the risk of infestation by the corresponding pathogen. However, the compounds of formula I can also penetrate the plant through the roots via the soil (systemic action) by drenching the locus of the plant with a liquid formulation, or by applying the compounds in solid form to the soil, e.g. in granular form (soil application). In crops of water rice such granulates can be applied to the flooded rice field. The compounds of formula I may also be applied to seeds (coating) by impregnating the seeds or tubers either with a liquid formulation of the fungicide or coating them with a solid formulation.

The formulation, i.e. the compositions containing the compound of formula I and, if desired, a solid or liquid adjuvant, are prepared in known manner, typically by intimately mixing and/or grinding the compound with extenders, e.g. solvents, solid carriers and, optionally, surface active compounds (surfactants).

The agrochemical formulations will usually contain from 0.1 to 99 % by weight, preferably from 0.1 to 95 % by weight, of the compound of formula I, 99.9 to 1 % by weight, preferably 99.8 to 5 % by weight, of a solid or liquid adjuvant, and from 0 to 25 % by weight, preferably from 0.1 to 25 % by weight, of a surfactant.

Advantageous rates of application are normally from 5 g to 2 kg of active ingredient (a.i.) per hectare (ha), preferably from 10 g to 1 kg a.i./ha, most preferably from 20 g to 600 g a.i./ha. When used as seed drenching agent, convenient dosages are from 10 mg to 1 g of active substance per kg of seeds.

Whereas it is preferred to formulate commercial products as concentrates, the end user will normally use dilute formulations.

The following non-limiting examples illustrate the above-described invention in more detail.


Temperatures are given in degrees Celsius. The following abbreviations are used:

m.p.= melting point; b.p.= boiling point. "NMR" means nuclear magnetic resonance spectrum.

MS stands for mass spectrum. "%" is percent by weight, unless corresponding concentrations are indicated in other units.

Example 1: 1-Methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [4-(4'-chlorophenyl)-pyridin-3-yl]amide

a) 1-Methyl-4-trifluoromethylpyrrole-3-carboxylic acid




Sodium hydride (8.0 g of a 75% dispersion in oil) is suspended at +5°C in a mixture of DMSO (300 ml) and diethylether (100 ml). A solution of ethyl 4,4,4-trifluorocrotonate (20 g) and TOSMIC (23 g) in DMSO (100 ml) is added through a dropping funnel at such a rate that the temperature does not exceed 10°C. After stirring the reaction mixture for an additional hour at room temperature methyl iodide (15.6 ml) is added with cooling. After 2 hours at room temperature the reaction mixture is poured onto crushed ice. Repeated extraction with ether, washing of the combined organic phases with brine and evaporation of the solvent under reduced pressure gives a product mixture in form of a light amber oil. The crude product mixture is heated at 60°C in a mixture of ethanol (100 ml) and sodium hydroxide (50 ml of a 30% aqueous solution). Washing of the solution with ether, acidifying of the aqueous phase with concentrated hydrochloric acid and filtering gives the 1-methyl-4-trifluoromethylpyrrole-3-

carboxylic acid in form of a crystalline solid. $^1\text{H-NMR}$ (CDCl_3): 7.24(d,1H); 6.88(d,1H); 3.63(s,3H).


In analogous manner the new compounds 1-ethyl- (m.p.146-148°C) and 1-methoxymethyl-4-trifluoromethylpyrrole-3-carboxylic acid can be prepared.

b) 4-(4'-chlorophenyl)-3-nitropyridine



In a sulfonation flask 6.9 g (44 mmol) of 4-chloro-3-nitropyridine, 6.5 g (42 mmol) 4-chlorophenylboronic acid, 6.0 g (44 mmol) potassium carbonate (saturated solution in water) and 1.0 g (0.9 mmol) tetrakis(triphenylphosphine)palladium are dissolved in 100 ml of dimethoxyethane (DME). The mixture is heated under reflux conditions under a constant nitrogen stream for 5 hours. Then the solvent is removed in a water jet vacuum and the residue taken up in ethylacetate. The organic phase is washed twice with water and after drying of the organic phase with sodium sulfate, the solvent is removed in a water jet vacuum. The resulting crude product is purified by column chromatography over silica gel (eluant: hexane/ethylacetate 1:1). Yield: 8.9 g 4-(4'-chlorophenyl)-3-nitropyridine in the form of brownish crystals; m.p. 74-76°C.


c) 3-amino-4-(4'-chlorophenyl)pyridine



In a sulfonation flask 8.4 g (36 mmol) 4-(4'-chlorophenyl)-3-nitropyridine is dissolved in a mixture of 100 ml water, 35 ml acetic acid and 10 ml n-propanol. After the addition of 7.0 g (125 mmol) iron powder, the mixture is heated for 3 hours under reflux conditions. After cooling to room temperature the mixture is diluted with ethylacetate and filtered over hyflo. Then the filtrate is neutralized by the addition of sodiumbicarbonate solution and the organic phase separated. The water phase is extracted twice with ethylacetate and the combined


organic phases dried over sodium sulfate. After distilling off the solvent in a water jet vacuum the obtained raw material is purified by column chromatography over silica gel (eluant: ethylacetate). Yield: 3.9 g 3-amino-4-(4'-chlorophenyl)pyridine in the form of a slightly brownish powder; m.p. : 144-146°C.

d) 1-Methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [4-(4'-chlorophenyl)-pyridin-3-yl]amide




A solution of 0.54 g (2.8 mmol) 1-methyl-4-trifluoromethylpyrrole-3-carboxylic acid and 0.39 g (3.1 mmol) oxalyl chloride in 20 ml methylene chloride is stirred for 2 hours at room temperature in the presence of a catalytic amount of DMF. Then the acid chloride solution is slowly added to a solution of 0.57 g (2.8 mmol) 3-amino-4-(4'-chlorophenyl)pyridine, 0.34 g (3.4 mmol) triethylamine and 15 ml methylene chloride. The resulting mixture is then stirred for 16 hours at room temperature. After the addition of ethylacetate, the organic phase is washed twice with water. After drying the organic phase over sodium sulfate, the solvent is removed in a water jet vacuum. The obtained crude product is purified by column chromatography over silica gel (eluant: hexane/THF 1:3). Yield: 0.4 g 1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [4-(4'-chlorophenyl)-pyridin-3-yl]amide in the form of yellowish crystals; m.p. : 178-180°C.

Example 2: 1-Methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid (2-bromo-benzamide)



A solution of 7.8 g (40.4 mmol) 1-methyl-4-trifluoromethylpyrrole-3-carboxylic acid and 3.8 ml (45 mmol) oxalyl chloride in 100 ml dichloromethane is stirred for 1 hour at room temperature in the presence of a catalytic amount of DMF. Then the resulting acid chloride solution is slowly added to a solution of 6.95 g (40.4 mmol) 2-bromoaniline, 6.9 ml (49 mmol) triethylamine and 140 ml dichloromethane. The resulting mixture is then stirred for 16 hours at room temperature. The organic phase is washed twice with water and dried over sodium sulfate. The solvent is removed under reduced pressure. The crude product is purified by column chromatography over silica gel (eluant: ethylacetate/hexane = 2/3); yield : 11.4 g 1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid (2-bromo-benzamide) as colourless crystals; m.p. 117-119°C.

Example 3: 1-Methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [2'-(5'-chloro-thienyl)-2-benzamide]




[cmpd. 1.20]

To 1.0 g (2.9 mmol) 1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid (2-bromo-benzamide), 0.56 g (3.5 mmol) 5-chloro-thiophene-2-boronic acid and 0.12 g (0.2 mmol) bis(triphenylphosphine)palladium(II) chloride in 20 ml ethylene glycol dimethyl ether is added a solution of 0.87 g (10.4 mmol) sodium hydrogencarbonate in 20 ml water. The reaction mixture is then stirred for 24 hours at 70°C. After the addition of dichloromethane, the organic phase is washed twice with water and with sodium chloride solution. After drying over sodium sulfate, the solvent is removed under reduced pressure. The product is purified by column chromatography over silica gel (eluant: dichloromethane/hexane = 4/1) to obtain 1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [2'-(5'-chloro-thienyl)-2-benzamide] as colourless crystals; m.p. 139-141°C.

















The following compounds are prepared in a similar way, using analogous methods.
















Table 1:
















Compounds of the formula I, wherein A = A1 =




















Compd. No.	R ₁	R ₂	X	R ₃	R ₄	Phys. Data [m.p. °C]
1.1	H	CH ₃	O		H	
1.2	H	CH ₃	O		H	
1.3	H	CH ₃	O		H	
1.4	H	CH ₃	O		H	
1.5	H	CH ₃	O		H	
1.6	H	CH ₃	O		H	
1.7	H	CH ₃	O		H	
1.8	H	CH ₃	O		H	
1.9	H	CH ₃	O		H	
1.10	H	CH ₃	O		H	
1.11	H	CH ₃	O		H	127-129
1.12	H	CH ₂ CH ₃	O		H	
1.13	H	CH ₃	S		H	

1.14	H	CH ₃	O		H	
1.15	H	CH ₃	O		H	131-132
1.16	H	CH ₃	S		H	resin
1.17	CH ₃	CH ₃	O		H	
1.18	H	CH ₂ OCH ₃	O		H	
1.19	H	CH ₃	O		H	116-118
1.20	H	CH ₃	O		H	139-141
1.21	H	CH ₃	O		H	132-136
1.22	H	CH ₃	O		H	
1.23	H	CH ₃	O		H	106-109
1.24	H	CH ₃	O		H	182-185
1.25	H	CH ₂ OCH ₃	O		H	171-173
1.26	H	CH ₃	O		H	132-134
1.27	H	CH ₂ OCH ₃	O		H	
1.28	H	CH ₃	O		H	
1.29	H	CH ₃	O		H	

1.30	H	CH ₃	O		H	
1.31	H	CH ₃	O		H	
1.32	H	CH ₃	O		H	
1.33	H	CH ₃	O		H	
1.34	H	CH ₂ CH ₃	O		H	
1.35	H	CH ₂ OCH ₃	O		H	
1.36	H	CH ₃	O		H	
1.37	H	CH ₃	O		H	
1.38	H	CH ₃	O		H	
1.39	H	CH ₃	O		H	
1.40	H	CH ₃	O		H	169-171
1.41	H	CH ₂ OCH ₃	O		H	
1.42	H	CH ₃	O		H	143-145
1.43	H	CH ₃	S		H	
1.44	H	CH ₃	O		H	

1.45	H	CH ₃	O		H	
1.46	H	CH ₂ OCH ₃	O		H	
1.47	H	CH ₃	O		H	
1.48	H	CH ₃	O		H	
1.49	H	CH ₃	O		H	> 230
1.50	H	CH ₂ OCH ₃	O		H	> 230
1.51	H	CH ₃	O		H	
1.52	H	CH ₃	O		H	
1.53	H	CH ₃	O		H	
1.54	H	CH ₃	O		H	
1.55	H	CH ₃	O		H	
1.56	H	CH ₃	O		H	
1.57	H	CH ₃	O		H	
1.58	H	CH ₃	O		H	
1.59	H	CH ₂ CH ₃	O		H	

1.60	H	CH ₃	O		H	
1.61	H	CH ₃	O		H	
1.62	H	CH ₃	O		H	
1.63	H	CH ₃	O		H	
1.64	H	CH ₃	O		H	
1.65	CH ₃	CH ₃	O		H	
1.66	H	CH ₃	O		H	126-128
1.67	H	CH ₃	O		H	164-165
1.68	H	CH ₃	O		H	118-119
1.69	H	CH ₂ OCH ₃	O		H	oil; ¹ H-NMR, MS
1.70	H	CH ₃	O		H	
1.71	H	CH ₂ OCH ₃	O		H	
1.72	H	CH ₃	O		H	
1.73	H	CH ₂ OCH ₃	O		H	
1.74	H	CH ₃	O		H	93-95
1.75	H	CH ₂ OCH ₃	O		H	oil; ¹ H-NMR, MS
1.76	H	CH ₃	O		H	













































1.77	H	CH ₂ OCH ₃	O		H	
1.78	H	CH ₃	O		H	
1.79	H	CH ₂ OCH ₃	O		H	
1.80	H	CH ₃	O		H	
1.81	H	CH ₃	O		H	250-252
1.82	H	CH ₃	O		H	202-204

Table 2: Compounds of the formula I, wherein A = A2 =


















Compd. No.	R ₁	R ₂	X	R ₃	R ₄	Phys. Data [m.p. °C]
2.1	H	CH ₃	O		H	
2.2	H	CH ₃	O		H	
2.3	H	CH ₃	O		H	
2.4	H	CH ₃	O		H	
2.5	H	CH ₃	O		H	
2.6	H	CH ₂ OCH ₃	O		H	

2.7	H	CH ₃	O		H	
2.8	H	CH ₃	O		H	
2.9	H	CH ₃	O		H	
2.10	H	CH ₃	O		H	
2.11	H	CH ₃	O		H	
2.12	H	CH ₃	O		H	resin
2.13	H	CH ₃	S		H	
2.14	H	CH ₃	O		H	
2.15	H	CH ₃	S		H	
2.16	H	CH ₃	O		H	
2.17	CH ₃	CH ₃	O		H	
2.18	H	CH ₂ OCH ₃	O		H	
2.19	H	CH ₃	O		H	
2.20	H	CH ₃	O		H	
2.21	H	CH ₃	O		H	

2.21	H	CH ₃	O		H	
2.22	H	CH ₃	O		H	
2.23	H	CH ₃	O		H	resin
2.24	H	CH ₂ OCH ₃	O		H	
2.25	H	CH ₃	O		H	
2.26	H	CH ₂ OCH ₃	O		H	
2.27	H	CH ₃	O		H	
2.28	H	CH ₃	O		H	
2.29	H	CH ₃	O		H	
2.30	H	CH ₃	O		H	
2.31	H	CH ₃	O		H	
2.32	CH ₃	CH ₃	O		H	
2.33	H	CH ₃	O		H	
2.34	H	CH ₂ CH ₃	O		H	
2.35	H	CH ₂ OCH ₃	O		H	
2.36	H	CH ₃	O		H	

-31 -

2.37	H	CH ₃	O		H	
2.38	H	CH ₃	O		H	
2.39	H	CH ₃	O		H	
2.40	H	CH ₃	O		H	resin
2.41	H	CH ₂ OCH ₃	O		H	
2.42	H	CH ₃	O		H	
2.43	H	CH ₃	S		H	
2.44	H	CH ₃	O		H	
2.45	H	CH ₃	O		H	
2.46	H	CH ₂ OCH ₃	O		H	
2.47	H	CH ₃	O		H	
2.48	H	CH ₃	O		H	
2.49	H	CH ₃	O		H	> 235
2.50	H	CH ₂ OCH ₃	O		H	
2.51	H	CH ₃	O		H	


















2.52	H	CH ₃	O		H
2.53	H	CH ₂ CH ₃	O		H
2.54	H	CH ₃	O		H
2.55	H	CH ₃	O		H
2.56	H	CH ₃	O		H
2.57	CH ₃	CH ₃	O		H
2.58	H	CH ₃	O		H
2.59	H	CH ₃	O		H
2.60	H	CH ₃	O		H
2.61	H	CH ₂ CH ₃	O		H
2.62	H	CH ₃	O		H
2.63	H	CH ₃	O		H
2.64	H	CH ₃	O		H
2.65	H	CH ₃	O		H
2.66	H	CH ₃	O		H
2.67	CH ₃	CH ₃	O		H

Table 3: Compounds of the formula I, wherein A = A17 =



Compd. No.	R ₁	R ₂	X	R ₃₁	R ₄	Phys. Data [m.p. °C]
3.1	H	CH ₃	O		H	
3.2	H	CH ₃	O		H	
3.3	H	CH ₃	O		H	
3.4	H	CH ₃	O		H	
3.5	H	CH ₂ OCH ₃	O		H	
3.6	H	CH ₃	O		H	
3.7	H	CH ₃	O		H	
3.8	H	CH ₃	O		H	
3.9	H	CH ₃	O		H	
3.10	H	CH ₃	O		H	
3.11	H	CH ₂ CH ₃	O		H	
















3.12	H	CH ₃	O		H
3.13	H	CH ₃	O		H
3.14	H	CH ₃	O		H
3.15	H	CH ₃	O		H
3.16	H	CH ₃	O		H
3.17	H	CH ₃	O		H
3.18	H	CH ₃	O		H
3.19	H	CH ₃	O		H
3.20	H	CH ₃	O		H
3.21	H	CH ₃	O		H
3.22	H	CH ₃	O		H
3.23	CH ₃	CH ₃	O		H
3.24	H	CH ₃	O		H
3.25	H	CH ₃	S		H

Table 4:

Compounds of the formula I, wherein A = A21 =



Compd. No.	R ₁	R ₂	X	R ₃₁	Phys. Data [m.p. °C]
4.1	H	CH ₃	O		
4.2	H	CH ₃	O		
4.3	H	CH ₃	O		
4.4	H	CH ₃	O		
4.5	H	CH ₂ OCH ₃	O		
4.6	H	CH ₃	O		
4.7	H	CH ₃	O		
4.8	H	CH ₃	O		
4.9	H	CH ₃	O		
4.10	H	CH ₃	O		resin
4.11	H	CH ₂ CH ₃	O		
4.12	H	CH ₃	O		















4.13	H	CH ₃	O	
4.14	H	CH ₃	O	
4.15	H	CH ₃	O	
4.16	H	CH ₃	O	
4.17	H	CH ₃	O	
4.18	H	CH ₃	O	
4.19	H	CH ₃	O	
4.20	H	CH ₃	O	
4.21	H	CH ₃	O	
4.22	H	CH ₃	O	
4.23	CH ₃	CH ₃	O	
4.24	H	CH ₃	O	
4.25	H	CH ₃	S	

Table 5: Compounds of the formula I, wherein A = A24 =



Compd. No.	R_1	R_2	X	R_{31}	R_4	Phys. Data [m.p. °C]
5.1	H	CH ₃	O		H	
5.2	H	CH ₃	O		H	
5.3	H	CH ₃	O		H	
5.4	H	CH ₃	O		H	
5.5	H	CH ₂ OCH ₃	O		H	
5.6	H	CH ₃	O		H	
5.7	H	CH ₃	O		H	
5.8	H	CH ₃	O		H	
5.9	H	CH ₃	O		H	
5.10	H	CH ₃	O		H	
5.11	H	CH ₂ CH ₃	O		H	
5.12	H	CH ₃	O		H	resin































5.13	H	CH ₃	O		H
5.14	H	CH ₃	O		H
5.15	H	CH ₃	O		H
5.16	H	CH ₃	O		H
5.17	H	CH ₃	O		H
5.18	H	CH ₃	O		H
5.19	H	CH ₃	O		H
5.20	H	CH ₃	O		H
5.21	H	CH ₃	O		H
5.22	H	CH ₃	O		H
5.23	CH ₃	CH ₃	O		H
5.24	H	CH ₃	O		H
5.25	H	CH ₃	S		H

Table 6: Compounds of formula I, wherein A = A25 =



Compd. No.	R_1	R_2	X	R_{31}	R_4	Phys. Data [m.p. °C]
6.1	H	CH ₃	O		H	
6.2	H	CH ₃	O		H	
6.3	H	CH ₃	O		H	
6.4	H	CH ₃	O		H	
6.5	H	CH ₂ OCH ₃	O		H	
6.6	H	CH ₃	O		H	
6.7	H	CH ₃	O		H	
6.8	H	CH ₃	O		H	
6.9	H	CH ₃	O		H	
6.10	H	CH ₃	O		H	178-180
6.11	H	CH ₂ CH ₃	O		H	
6.12	H	CH ₂ OCH ₃	O		H	

6.13	H	CH ₃	O		H
6.14	H	CH ₃	O		H
6.15	H	CH ₃	O		H
6.16	H	CH ₃	S		H
6.17	H	CH ₃	O		H
6.18	H	CH ₃	O		H
6.19	H	CH ₃	O		H
6.20	H	CH ₃	O		H
6.21	H	CH ₃	O		H
6.22	H	CH ₃	O		H
6.23	H	CH ₃	O		H
6.24	H	CH ₃	O		H
6.25	H	CH ₃	O		H
6.26	H	CH ₃	O		H
6.27	H	CH ₃	O		H
6.28	H	CH ₃	O		H

165-167









6.29	H	CH ₃	O		H
6.30	H	CH ₃	O		H
6.31	CH ₃	CH ₃	O		H
6.32	H	CH ₃	O		H
6.33	H	CH ₃	S		H
6.34	H	CH ₃	O		H
6.35	H	CH ₃	O		H

Table 7: Compounds of the formula I, wherein A = A26 =



Compd. No.	R ₁	R ₂	X	R ₃₁	R ₄	phys. data m.p. °C
7.1	H	CH ₃	O		H	
7.2	H	CH ₃	O		H	resin
7.3	H	CH ₂ CH ₃	O		H	
7.4	H	CH ₂ OCH ₃	O		H	
7.5	H	CH ₃	O		H	
7.6	H	CH ₂ OCH ₃	O		H	
7.7	H	CH ₃	S		H	
7.8	H	CH ₃	O		H	
7.9	H	CH ₃	O		H	
7.10	CH ₃	CH ₃	O		H	
7.11	H	CH ₃	O		H	
7.12	H	CH ₃	O		H	













7.13	H	CH ₃	O		H
7.14	H	CH ₃	O		H
7.15	H	CH ₃	O		H

Table 8:

Compounds of the formula I, wherein A = A27 =



Compd. No.	R ₁	R ₂	X	R ₃₁	R ₄	phys. data m.p. °C
8.1	H	CH ₃	O		H	63-65
8.2	H	CH ₃	O		H	156-157
8.3	H	CH ₂ CH ₃	O		H	
8.4	H	CH ₂ OCH ₃	O		H	
8.5	H	CH ₃	O		H	119-122
8.6	H	CH ₂ OCH ₃	O		H	
8.7	H	CH ₃	S		H	
8.8	H	CH ₃	O		H	












8.9	H	CH ₃	O		H	
8.10	CH ₃	CH ₃	O		H	
8.11	H	CH ₃	O		H	
8.12	H	CH ₃	O		H	
8.13	H	CH ₃	O		H	
8.14	H	CH ₃	O		H	159-161
8.15	H	CH ₂ OCH ₃	O		H	139-141
8.16	H	CH ₃	O		H	140-143
8.17	H	CH ₃	O		H	
8.18	H	CH ₃	O		H	

Table 9:


Compounds of the formula I, wherein A = A31 =

















Compd. No.	R ₁	R ₂	X	R ₄	R ₅	R ₆	R ₇	R ₈	phys.data m.p. °C
9.1	H	CH ₃	O	H	H	CH ₃	CH ₃	CH ₃	165-167
9.2	H	CH ₂ CH ₃	O	H	H	CH ₃	CH ₃	CH ₃	












9.3	H	CH ₂ OCH ₃	O	H	H	CH ₃	CH ₃	CH ₃	oil; MS
9.4	CH ₃	CH ₃	O	H	H	CH ₃	CH ₃	CH ₃	
9.5	H	CH ₃	S	H	H	CH ₃	CH ₃	CH ₃	
9.6	H	CH ₃	O	H	CH ₃	CH ₃	CH ₃	H	
9.7	H	CH ₂ OCH ₃	O	H	CH ₃	CH ₃	CH ₃	H	

Table 10: Compounds of formula I, wherein A = A22 =



Compd. No.	R ₁	R ₂	X	R ₃₁	R ₄	Phys. Data [m.p. °C]
10.1	H	CH ₃	O		H	
10.2	H	CH ₃	O		H	
10.3	H	CH ₃	O		H	oil
10.4	H	CH ₃	O		H	
10.5	H	CH ₂ OCH ₃	O		H	
10.6	H	CH ₃	O		H	
10.7	H	CH ₃	O		H	
10.8	H	CH ₃	O		H	

10.9	H	CH ₃	O		H	
10.10	H	CH ₃	O		H	
10.11	H	CH ₂ CH ₃	O		H	
10.12	H	CH ₂ OCH ₃	O		H	
10.13	H	CH ₃	O		H	
10.14	H	CH ₃	O		H	
10.15	H	CH ₃	O		H	216-219
10.16	H	CH ₃	S		H	
10.17	H	CH ₃	O		H	
10.18	H	CH ₃	O		H	
10.19	H	CH ₃	O		H	
10.20	H	CH ₃	O		H	
10.21	H	CH ₃	O		H	
10.22	H	CH ₃	O		H	
10.23	H	CH ₃	O		H	
10.24	H	CH ₃	O		H	

10.25	H	CH ₃	O		H
10.26	H	CH ₃	O		H
10.27	H	CH ₃	O		H
10.28	H	CH ₃	O		H
10.29	H	CH ₃	O		H
10.30	H	CH ₃	O		H
10.31	CH ₃	CH ₃	O		H
10.32	H	CH ₃	O		H
10.33	H	CH ₃	S		H
10.34	H	CH ₃	O		H
10.35	H	CH ₃	O		H

Formulation Examples for compounds of formula I

Working procedures for preparing formulations of the compounds of formula I such as Emulsifiable concentrates, Solutions, Granulates, Dusts and Wettable powders are described in WO 97/33890.

Biological Examples: Fungicidal actionsExample B-1: Action against *Puccinia recondita* / wheat (Brownrust on wheat)

1 week old wheat plants cv. Arina are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application wheat plants are inoculated by spraying a spore suspension (1×10^5 uredospores/ml) on the test plants. After an incubation period of 2 days at 20° C and 95% r. h. plants are kept in a greenhouse for 8 days at 20° C and 60% r.h. The disease incidence is assessed 10 days after inoculation. Compounds of Tables 1 to 10 show good activity in these tests. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

Example B-2: Action against *Podosphaera leucotricha* / apple (Powdery mildew on apple)

5 week old apple seedlings cv. McIntosh are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. One day after application apple plants are inoculated by shaking plants infected with apple powdery mildew above the test plants. After an incubation period of 12 days at 22° C and 60% r. h. under a light regime of 14/10 h (light/dark) the disease incidence is assessed.

Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

Example B-3: Action against *Venturia inaequalis* / apple (Scab on apple)

4 week old apple seedlings cv. McIntosh are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application apple plants are inoculated by spraying a spore suspension (4×10^5 conidia/ml) on the test plants. After an incubation period of 4 days at 21° C and 95% r. h. the plants are placed for 4 days at 21° C

and 60% r. h. in a greenhouse. After another 4 day incubation period at 21° C and 95% r. h. the disease incidence is assessed.

Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

Example B-4: Action against *Erysiphe graminis* / barley (Powdery mildew on barley)

1 week old barley plants cv. Express are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application barley plants are inoculated by shaking powdery mildew infected plants above the test plants. After an incubation period of 6 days at 20° C / 18° C (day/night) and 60% r. h. in a greenhouse the disease incidence is assessed.

Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

Example B-5: Action against *Botrytis cinerea* / apple (Botrytis on apple fruits)

In an apple fruit cv. Golden Delicious 3 holes are drilled and each filled with 30 µl droplets of the formulated test compound (0.002% active ingredient). Two hours after application 50 µl of a spore suspension of *B. cinerea* (4×10^5 conidia/ml) are pipetted on the application sites. After an incubation period of 7 days at 22° C in a growth chamber the disease incidence is assessed.

Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

Example B-6: Action against *Botrytis cinerea* / grape (Botrytis on grapes)

5 week old grape seedlings cv. Gutedel are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. Two days after application grape plants are inoculated by spraying a spore suspension (1×10^6 conidia/ml) on the test plants. After an

incubation period of 4 days at 21° C and 95% r. h. in a greenhouse the disease incidence is assessed.

Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

Example B-7: Action against *Botrytis cinerea* / tomato (Botrytis on tomatoes)

4 week old tomato plants cv. Roter Gnom are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. Two days after application tomato plants are inoculated by spraying a spore suspension (1×10^5 conidia/ml) on the test plants. After an incubation period of 4 days at 20° C and 95% r. h. in a growth chamber the disease incidence is assessed.

Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

Example B-8: Action against *Pyrenophora teres* / barley (Net blotch on barley)

1 week old barley plants cv. Express are treated with the formulated test compound (0.002% active ingredient) in a spray chamber. Two days after application barley plants are inoculated by spraying a spore suspension (3×10^4 conidia/ml) on the test plants. After an incubation period of 2 days at 20° C and 95% r. h. plants are kept for 2 days at 20° C and 60% r.h. in a greenhouse. The disease incidence is assessed 4 days after inoculation.

Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).


Example B-9: Action against *Septoria nodorum* / wheat (Septoria leaf spot on wheat)

1 week old wheat plants cv. Arina are treated with the formulated test compound (0.02% active ingredient) in a spray chamber. One day after application wheat plants are inoculated by spraying a spore suspension (5×10^5 conidia/ml) on the test plants. After an

incubation period of 1 day at 20° C and 95% r. h. plants are kept for 10 days at 20° C and 60% r.h. in a greenhouse. The disease incidence is assessed 11 days after inoculation. Compounds of Tables 1 to 10 show good activity in this test. The compounds 1.11, 1.15, 1.16, 1.19-1.21, 1.23-1.26, 1.40, 1.42, 1.49, 1.66-1.69, 1.74, 1.75, 1.81, 2.12, 2.23, 2.40, 2.49, 4.10, 5.12, 6.10, 6.15, 7.2, 8.1, 8.2, 8.5, 8.14-8.16, 10.3 and 10.15 exhibit strong efficacy (< 20% infestation).

What is claimed is

1. A trifluoromethylpyrrolcarboxamide of the formula I



wherein

X is oxygen or sulfur;

R₁ is hydrogen, C₁-C₄alkyl unsubstituted or substituted, or halogen;

R₂ is C₁-C₄alkyl unsubstituted or substituted; and
















A is orthosubstituted aryl; orthosubstituted heteroaryl; bicycloaryl unsubstituted or substituted; or bicycloheteroaryl unsubstituted or substituted.


2. A compound of formula I according to claim 1, wherein

R₁ is hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄haloalkoxy-C₁-C₄alkyl or halogen ;

R₂ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy-C₁-C₄alkyl or C₁-C₄haloalkoxy-C₁-C₄alkyl ;

A is a group





and

R_3 is C_3 - C_7 cycloalkyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkoxy or C_1 - C_4 alkyl; C_4 - C_7 cycloalkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkoxy or C_1 - C_4 alkyl; C_6 - C_7 cyclodialkenyl unsubstituted or mono- to trisubstituted by halogen, hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 haloalkoxy or C_1 - C_4 alkyl; thienyl, furyl, pyrrolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, thiadiazolyl, imidazolyl, triazinyl, benzothienyl, tetrazolyl, 5,6-dihydro-1,4,2-dioxazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl, which are unsubstituted or substituted by halogen, C_1 - C_6 haloalkyl, C_1 - C_6 alkyl, hydroxy, cyano, nitro, CHO, C_1 - C_4 alkoxy- C_1 - C_4 alkyl, COOC- C_4 alkyl, C_1 - C_6 haloalkoxy- C_1 - C_4 alkyl, C_1 - C_6 alkoxy or C_1 - C_6 haloalkoxy;

R_{31} is C_3 - C_7 cycloalkyl, C_3 - C_7 cycloalkenyl or C_5 - C_7 cycloalkadienyl which are unsubstituted or substituted by halogen, C_1 - C_6 alkoxy, C_1 - C_6 alkoxy- C_1 - C_6 alkyl, C_1 - C_6 haloalkoxy- C_1 - C_4 alkyl,

C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄haloalkoxy, C₂-C₄alkenyl or C₂-C₅alkynyl; phenyl unsubstituted or substituted by halogen, C₁-C₆alkoxy, C₁-C₆haloalkoxy, C₁-C₄alkyl, C₁-C₄haloalkyl, C₂-C₄alkenyl, C₂-C₅alkynyl, CHO, COOC₁-C₄alkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₄alkyl-C₁-C₄alkoxy, C₁-C₄haloalkoxy-C₁-C₄alkyl, C₁-C₄haloalkyl-C₁-C₄alkoxy, cyano or nitro; thienyl, furyl, pyrrolyl, pyrazolyl, oxazolyl, thiazolyl, isoxazolyl, isothiazolyl, thiadiazolyl, imidazolyl, triazinyl, pyridyl, pyridazinyl, pyrazinyl or pyrimidinyl, which are unsubstituted or substituted by halogen, C₁-C₆haloalkyl, C₁-C₆alkyl, hydroxy, cyano, nitro, CHO, COOC₁-C₄alkyl, C₁-C₆haloalkoxy-C₁-C₄alkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₆alkoxy or C₁-C₆haloalkoxy;

R₄ is hydrogen; cyano; nitro; halogen; C₁-C₄alkoxy; C₁-C₄haloalkyl; C₁-C₄alkyl; C₁-C₄alkoxy-C₁-C₄alkyl; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₄haloalkoxy-C₁-C₄alkyl; or C₁-C₄haloalkoxy; and

R₅, R₆, R₇, R₈ and R₉ are identical or different and are each independently of the others hydrogen, halogen, C₁-C₄haloalkyl, C₁-C₄alkyl, C₁-C₆haloalkoxy-C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkoxy, C₁-C₄alkoxy-C₁-C₄alkyl or C₃-C₇cycloalkyl.

3. A compound of formula I according to claim 2, wherein

X is oxygen.

4. A compound of formula I according to claim 2, wherein

X is sulfur.

5. A compound of formula I according to claim 3, wherein

R₁ is hydrogen, C₁-C₄alkyl or C₁-C₄haloalkyl;

R₂ is C₁-C₄alkyl, C₁-C₃haloalkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

A is A1, A2, A3, A5, A8, A10, A13, A14, A17, A18, A20, A21, A22, A24, A25, A26, A27, A29, A31 or A32;

R₃ is C₅-C₇cycloalkyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₅-C₇cycloalkenyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₆-C₇cyclodialkenyl, unsubstituted or mono- to disubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl, which are

unsubstituted or substituted by halogen, hydroxy, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy or C₁-C₄haloalkoxy;

R₃₁ is C₅-C₇cycloalkyl, C₅-C₇cycloalkenyl or C₅-C₇cycloalkadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; phenyl which is unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₄ is hydrogen; halogen; C₁-C₄alkyl; C₁-C₄alkoxy; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₄haloalkyl; or C₁-C₄haloalkoxy; and

R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy.

6. A compound of formula I according to claim 5, wherein

A is A1, A2, A3, A17, A20, A21, A24, A25, A26, A27 or A31;

R₁ is hydrogen or C₁-C₃alkyl;

R₂ is C₁-C₃alkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

R₃ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or mono- to disubstituted by chloro, bromo, C₁-C₂alkyl, C₁-C₂haloalkyl or C₁-C₂haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₃₁ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₄ is hydrogen; halogen; C₁-C₃alkyl; C₁-C₃haloalkyl; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₃alkoxy; or C₁-C₃haloalkoxy; and

R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen or C₁-C₃alkyl.

7. A compound of formula I according to claim 4, wherein

R₁ is hydrogen, C₁-C₄alkyl or C₁-C₄haloalkyl;

R₂ is C₁-C₄alkyl, C₁-C₃haloalkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

A is A1, A2, A3, A5, A8, A10, A13, A14, A17, A18, A20, A21, A22, A24, A25, A26, A27, A29, A31 or A32;

R₃ is C₅-C₇cycloalkyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₅-C₇cycloalkenyl, unsubstituted or mono- to trisubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; C₆-C₇cyclodialkenyl, unsubstituted or mono- to disubstituted by halogen, hydroxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₃alkyl or C₁-C₃alkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl, which are unsubstituted or substituted by halogen, hydroxy, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy or C₁-C₄haloalkoxy;

R₃₁ is C₅-C₇cycloalkyl, C₅-C₇cycloalkenyl or C₅-C₇cycloakadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₂-C₄alkenyl, C₂-C₄alkynyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; phenyl which is unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, isoxazolyl, oxazolyl, thiadiazolyl, triazinyl, pyridyl, pyrimidinyl, pyrazinyl or pyridazinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₄ is hydrogen; halogen; C₁-C₄alkyl; C₁-C₄alkoxy; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₄haloalkyl; or C₁-C₄haloalkoxy; and

R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy.

8. A compound of formula I according to claim 7, wherein

A is A1, A2, A3, A17, A20, A21, A24, A25, A26, A27 or A31;

R₁ is hydrogen or C₁-C₃alkyl;

R₂ is C₁-C₃alkyl or C₁-C₃alkoxy-C₁-C₃alkyl;

R₃ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or mono- to disubstituted by chloro, bromo, C₁-C₂alkyl, C₁-C₂haloalkyl or C₁-C₂haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₃₁ is cyclohexyl, cyclohexenyl or cyclohexadienyl, which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄alkoxy, C₁-C₄haloalkyl or C₁-C₄haloalkoxy; thienyl, furyl, triazinyl, pyridyl, pyrazinyl, pyridazinyl or pyrimidinyl which are unsubstituted or substituted by halogen, C₁-C₄alkyl, C₁-C₄haloalkyl or C₁-C₄haloalkoxy;

R₄ is hydrogen; halogen; C₁-C₃alkyl; C₁-C₃haloalkyl; C₅-C₇cycloalkyl unsubstituted or substituted by C₁-C₃alkyl or C₁-C₃haloalkyl; C₁-C₃alkoxy; or C₁-C₃haloalkoxy; and R₅, R₆, R₇ and R₈ are identical or different and are each independently of the others hydrogen or C₁-C₃alkyl.

9. A compound of formula I according to claim 3, wherein

A is A1 and

R₃ is C₅-C₇cycloalkyl unsubstituted or monosubstituted by C₁-C₄alkyl or C₁-C₄haloalkyl.

10. A compound of formula I according to claim 9 selected from the group comprising

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-methylcyclohexyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-ethylcyclohexyl)-2-benzamide];


1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-trifluoromethylcyclohexyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-methylcyclopentyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-ethylcyclopentyl)-2-benzamide];

1-methyl-4-trifluoromethyl-1H-pyrrole-3-carboxylic acid [1'-(3'-trifluoromethylcyclopentyl)-2-benzamide].


11. A process for the preparation of compounds of formula I which comprises reacting the starting materials according to the scheme



Base = NEt_3 , Hünig-base, Na_2CO_3 , K_2CO_3 and others

wherein R_1 , R_2 and A are as defined for formula I in claim 1;

or when A is phenyl, pyridine or pyrimidine according to the scheme



G = phenyl, pyridine, pyrimidine

wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1.


12. A composition for controlling microorganisms and preventing attack and infestation of plants therewith, wherein the active ingredient is a compound as claimed in claim 1 together with a suitable carrier.

13. Use of a compound of formula I according to claim 1 for protecting plants against infestation by phytopathogenic microorganisms.

14. A method of controlling or preventing infestation of cultivated plants by phytopathogenic microorganisms by application of a compound of formula I as claimed in claim 1 to plants, to parts thereof or the locus thereof.

[received by the International Bureau on 02 March 2001 (02.03.01);
original claim 1 amended; remaining claims unchanged (1 page)]

1. A trifluoromethylpyrrolcarboxamide of the formula I




wherein

X is oxygen or sulfur;

R₁ is hydrogen, C₁-C₄alkyl unsubstituted or substituted, or halogen;

R₂ is C₁-C₄alkyl unsubstituted or substituted; and

A is orthosubstituted aryl; orthosubstituted heteroaryl; bicycloaryl unsubstituted or substituted; or bicycloheteroaryl unsubstituted or substituted; with the proviso that when X=O



C₃-C₇cycloalkyl or unsubstituted C₄-C₇cycloalkenyl.


2. A compound of formula I according to claim 1, wherein

R₁ is hydrogen, C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy-C₁-C₄alkyl, C₁-C₄alkoxy,

C₁-C₄haloalkoxy, C₁-C₄haloalkoxy-C₁-C₄alkyl or halogen ;

R₂ is C₁-C₄alkyl, C₁-C₄haloalkyl, C₁-C₄alkoxy-C₁-C₄alkyl or C₁-C₄haloalkoxy-C₁-C₄alkyl ;

A is a group



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/11196

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D207/34 C07D401/12 C07D409/14 C07D407/12 C07D413/12
 C07D409/12 A01N43/36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 00 09482 A (NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT) 24 February 2000 (2000-02-24) claims 1-12	1-14
A	US 5 162 308 A (D. G. BROWN ET AL.) 10 November 1992 (1992-11-10) claims 1-13	1-14
A	EP 0 372 263 A (AMERICAN CYANAMID CO.) 13 June 1990 (1990-06-13) claims 1-21	1-14
A	EP 0 481 182 A (AMERICAN CYANAMID CO.) 22 April 1992 (1992-04-22) claims 1-12	1-14



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

19 February 2001

Date of mailing of the international search report

26/02/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Herz, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/11196

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 600 157 A (AMERICAN CYANAMID CO.) 8 June 1994 (1994-06-08) claims 1-10 ---	1-14
A	WO 97 08148 A (BASF AG) 6 March 1997 (1997-03-06) claims 1-6 ---	1-14
A	EP 0 737 682 A (MITSUI TOATSU CHEMCALS, INC.) 16 October 1996 (1996-10-16) claims 1-17 ---	1-14
A	EP 0 566 138 A (HODOGAYA CHEMICAL CO., LTD.) 20 October 1993 (1993-10-20) claims 1-4 -----	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/EP 00/11196

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0009482	A	24-02-2000	AU 5513899 A	06-03-2000
US 5162308	A	10-11-1992	US 5204332 A	20-04-1993
			AT 118303 T	15-03-1995
			AU 625743 B	16-07-1992
			AU 4586689 A	07-06-1990
			BR 8906202 A	25-09-1990
			CA 2004388 A	05-06-1990
			CN 1043317 A,B	27-06-1990
			CN 1098590 A,B	15-02-1995
			CZ 8906825 A	14-07-1999
			DD 289694 A	08-05-1991
			DE 68921160 D	23-03-1995
			DE 68921160 T	19-10-1995
			DK 609589 A	06-06-1990
			EP 0372263 A	13-06-1990
			ES 2068229 T	16-04-1995
			FI 97293 B	15-08-1996
			GR 3015061 T	31-05-1995
			HU 53485 A,B	28-11-1990
			IE 67281 B	20-03-1996
			IL 92311 A	19-01-1996
			IN 170306 A	14-03-1992
			JP 2237971 A	20-09-1990
			JP 2888568 B	10-05-1999
			KR 136105 B	25-04-1998
			NZ 231568 A	27-01-1993
			RU 2109012 C	20-04-1998
			RU 2011346 C	30-04-1994
			ZA 8909252 A	26-09-1990
EP 372263	A	13-06-1990	AT 118303 T	15-03-1995
			AU 625743 B	16-07-1992
			AU 4586689 A	07-06-1990
			BR 8906202 A	25-09-1990
			CA 2004388 A	05-06-1990
			CN 1043317 A,B	27-06-1990
			CN 1098590 A,B	15-02-1995
			CZ 8906825 A	14-07-1999
			DD 289694 A	08-05-1991
			DE 68921160 D	23-03-1995
			DE 68921160 T	19-10-1995
			DK 609589 A	06-06-1990
			ES 2068229 T	16-04-1995
			FI 97293 B	15-08-1996
			GR 3015061 T	31-05-1995
			HU 53485 A,B	28-11-1990
			IE 67281 B	20-03-1996
			IL 92311 A	19-01-1996
			IN 170306 A	14-03-1992
			JP 2237971 A	20-09-1990
			JP 2888568 B	10-05-1999
			KR 136105 B	25-04-1998
			NZ 231568 A	27-01-1993
			RU 2109012 C	20-04-1998
			RU 2011346 C	30-04-1994
			US 5162308 A	10-11-1992
			US 5204332 A	20-04-1993

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP 00/11196

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 372263 A		ZA 8909252 A	26-09-1990
EP 481182 A	22-04-1992	US 5157047 A	20-10-1992
		AT 148455 T	15-02-1997
		AU 646693 B	03-03-1994
		AU 8595391 A	30-04-1992
		BR 9104517 A	09-06-1992
		CA 2053513 A	19-04-1992
		DE 69124452 D	13-03-1997
		DE 69124452 T	15-05-1997
		DK 481182 T	17-02-1997
		ES 2096604 T	16-03-1997
		GR 3022451 T	30-04-1997
		IL 99209 A	14-05-1996
		JP 4288054 A	13-10-1992
		PT 99242 A, B	30-09-1992
		US 5281719 A	25-01-1994
		US 5229524 A	20-07-1993
		ZA 9108295 A	29-07-1992
EP 600157 A	08-06-1994	US 5449789 A	12-09-1995
		AU 5201293 A	09-06-1994
		BR 9304870 A	14-06-1994
		CA 2110089 A	31-05-1994
		CN 1091127 A	24-08-1994
		CZ 9302426 A	15-06-1994
		HU 65577 A, B	28-07-1994
		JP 6247930 A	06-09-1994
		MX 9306604 A	31-05-1994
		SK 129193 A	09-11-1994
		US 5563279 A	08-10-1996
		ZA 9308919 A	02-08-1994
WO 9708148 A	06-03-1997	DE 19531813 A	06-03-1997
		AU 6928596 A	19-03-1997
		EP 0847388 A	17-06-1998
		JP 11511449 T	05-10-1999
		US 5998450 A	07-12-1999
EP 737682 A	16-10-1996	CA 2173788 A	12-10-1996
		CN 1146993 A, B	09-04-1997
		JP 9235282 A	09-09-1997
		KR 201426 B	15-06-1999
		US 5747518 A	05-05-1998
EP 566138 A	20-10-1993	DE 69322118 D	24-12-1998
		DE 69322118 T	20-05-1999
		JP 6025199 A	01-02-1994